
Flames of Relief: Fire Cupping Therapy for Lumbo-Sacral Pain Management - A Case Study Analysis of Five Patients

*Izharul Hasan**

**Consultant Unani, Ayush Wellness Centre, President's Estate, New Delhi*

**Corresponding Author*

Email: drizharnium@gmail.com

ABSTRACT

This case study explores the efficacy of fire cupping therapy in managing lumbo-sacral pain through the analysis of five patient cases. The study aims to investigate the outcomes of fire cupping therapy sessions in alleviating pain and improving the quality of life for individuals suffering from lumbo-sacral pain. The methodology involved selecting patients based on specific criteria, administering fire cupping therapy sessions, and conducting follow-up assessments to evaluate treatment responses. Results indicate significant reductions in pain levels across all patients following fire cupping therapy, with minimal adverse effects observed. The findings suggest that fire cupping therapy holds promise as a complementary approach for lumbo-sacral pain management. This study underscores the potential of fire cupping therapy in providing relief for individuals with lumbo-sacral pain and emphasizes the need for further research to elucidate its mechanisms of action and optimize its clinical application.

Keywords: *Fire cupping therapy, Lumbo-sacral pain, Pain management*

INTRODUCTION

Lumbo-sacral pain, affecting the lower back and sacral region, is a common musculoskeletal complaint with various etiologies, including degenerative conditions, muscle strains, and nerve impingement. It often leads to significant discomfort, functional limitations, and reduced quality of life for affected individuals [1]. Lower back pain is one of the most prevalent musculoskeletal disorders worldwide, affecting people of all ages and socioeconomic backgrounds. Epidemiological studies indicate that the lifetime prevalence of lumbo-sacral pain ranges from 60% to 80%, with the point prevalence estimated to be around 15% to 30% at any given time.

Lumbo-sacral pain imposes a significant socioeconomic burden on healthcare systems, employers, and individuals. It ranks among the leading causes of disability-adjusted life years (DALYs) globally, contributing to lost productivity, work absenteeism, and healthcare expenditures. Fire cupping therapy is an ancient healing modality that involves placing heated glass or bamboo cups on the skin to create a vacuum, drawing underlying tissues into the cup. This practice is rooted in traditional Chinese medicine as well as in Unani medicine and is believed to promote blood circulation, release muscle tension, and alleviate pain. Fire cupping therapy has gained popularity as a complementary and alternative treatment for various musculoskeletal conditions, including lumbo-sacral pain [2].

The primary aim of this case study is to investigate the effectiveness of fire cupping therapy in managing lumbo-sacral pain. By examining the experiences of five patients who underwent fire cupping therapy, this study seeks to evaluate the outcomes of treatment,

including changes in pain levels, functional improvements, and any adverse effects. Through comprehensive analysis and discussion, this case study aims to contribute to the existing literature on fire cupping therapy and provide insights into its potential role in lumbo-sacral pain management.

LITERATURE REVIEW

Lumbo-sacral pain management encompasses a variety of treatment modalities, including pharmacological interventions, physical therapy, exercise programs, and complementary therapies. These approaches aim to alleviate pain, improve function, and enhance the overall well-being of individuals affected by lumbo-sacral pain. While some treatments focus on symptom relief, others target underlying pathology to address the root cause of pain [3].

Several studies have investigated the use of fire cupping therapy for pain management, including its application in conditions such as back pain, neck pain, and musculoskeletal disorders. These studies have employed diverse methodologies, including randomized controlled trials, observational studies, and case reports, to evaluate the efficacy and safety of fire cupping therapy. Findings from these studies have suggested potential benefits of fire cupping therapy in reducing pain intensity, improving range of motion, and enhancing quality of life for patients with various pain conditions [4, 5].

The existing evidence on the effectiveness of fire cupping therapy for pain management is mixed, with some studies reporting positive outcomes while others yield inconclusive results. While proponents of fire cupping therapy emphasize its ability to stimulate blood flow, release muscle tension, and promote healing, skeptics question its mechanisms of action and suggest that any perceived benefits may be attributed to a placebo effect or other factors. Despite the ongoing debate, there is growing interest in further exploring the therapeutic potential of fire cupping therapy and elucidating its role in pain management paradigms [6, 7].

METHODOLOGY

A. Selection Criteria for Patients: The selection criteria for patients participating in this case study include individuals aged 18 years or older experiencing chronic or recurrent lumbo-sacral pain of at least moderate intensity (as assessed by a numerical rating scale). Patients with a history of previous treatments for lumbo-sacral pain, such as medication, physical therapy, or surgery, are eligible for inclusion. Exclusion criteria comprise individuals with contraindications to fire cupping therapy, such as skin infections, bleeding disorders, or pregnancy, as well as those unable to provide informed consent or comply with study requirements [8].

B. Description of Fire Cupping Procedure: The fire cupping procedure involves the following steps:

- 1) Pre-treatment assessment: Patients undergo a thorough evaluation of their medical history, current symptoms, and pain intensity.
- 2) Preparation: The treatment area is cleansed, and the patient is positioned comfortably in a prone or supine position.
- 3) Application of lubricant: Roghan Baboona, is applied to the skin to facilitate movement of the cups and prevent skin irritation.
- 4) Heating of cups: Glass cups are heated using an open flame or other heating source until the air inside the cup heats up and creates a vacuum.

- 5) Cup placement: The heated cups are quickly placed onto specific acupoints or areas of tension along the lumbo-sacral region, creating suction and drawing underlying tissues into the cups.
- 6) Duration of treatment: The cups are left in place for a predetermined period, typically ranging from 5 to 15 minutes, depending on patient tolerance and therapist discretion.
- 7) Removal of cups: After the designated treatment duration, the cups are carefully removed, and the treatment area is assessed for any adverse reactions.
- 8) Post-treatment care: Patients may receive post-treatment instructions, such as avoiding exposure to cold or drafts and staying hydrated [9].

C. Data Collection Methods: Data collection includes both quantitative and qualitative measures:

- 1) Quantitative measures: Pain intensity scores, functional assessments (e.g., Oswestry Disability Index), and any adverse events are recorded before and after each fire cupping session.
- 2) Qualitative measures: Patients' subjective experiences, including changes in pain perception, mobility, and overall well-being, are documented through structured interviews or patient-reported outcome measures.

CASE PRESENTATION

A. Patient 1:

- Demographic information: Male, 45 years old.
- Medical history: History of chronic low back pain due to degenerative disc disease. Previous treatments include physical therapy and nonsteroidal anti-inflammatory drugs (NSAIDs).
- Presenting symptoms: Patient reports persistent dull, aching pain in the lumbo-sacral region, aggravated by prolonged sitting and standing. Pain intensity rated at 7/10 on the numerical rating scale.

B. Patient 2:

- Demographic information: Female, 55 years old.
- Medical history: History of lumbar spinal stenosis and osteoarthritis. Underwent lumbar laminectomy surgery two years ago. Current treatments include acetaminophen and occasional use of muscle relaxants.
- Presenting symptoms: Patient complains of sharp, shooting pain radiating from the lower back to the buttocks and thighs, exacerbated by walking and prolonged standing. Pain intensity rated at 8/10 on the numerical rating scale.

C. Patient 3:

- Demographic information: Male, 60 years old.
- Medical history: History of work-related lumbar strain. Underwent physical therapy and chiropractic treatments in the past.
- Presenting symptoms: Patient describes aching and stiffness in the lower back, worsened by bending and lifting activities. Reports difficulty in performing daily tasks due to pain. Pain intensity rated at 6/10 on the numerical rating scale.

D. Patient 4:

- Demographic information: Female, 40 years old.

- Medical history: History of fibromyalgia and irritable bowel syndrome. Currently managed with medications, including tricyclic antidepressants and selective serotonin reuptake inhibitors (SSRIs).
- Presenting symptoms: Patient experiences diffuse, chronic pain involving the lower back and pelvis, associated with generalized muscle stiffness and fatigue. Pain intensity rated at 6/10 on the numerical rating scale.

E. Patient 5:

- Demographic information: Male, 50 years old.
- Medical history: History of chronic low back pain following a work-related injury. Underwent multiple epidural steroid injections and facet joint injections in the past.
- Presenting symptoms: Patient reports persistent, stabbing pain localized to the lumbar spine, exacerbated by bending and twisting motions. Pain intensity rated at 7/10 on the numerical rating scale.

Table 1: Patient Characteristics

Patient	Age (years)	Gender	Medical History	Presenting Symptoms
1	45	Male	Chronic low back pain due to degenerative disc disease	Dull, aching pain in lumbo-sacral region, aggravated by prolonged sitting and standing
2	55	Female	Lumbar spinal stenosis, osteoarthritis, previous lumbar laminectomy surgery	Sharp, shooting pain radiating from lower back to buttocks and thighs, exacerbated by walking and prolonged standing
3	60	Male	Work-related lumbar strain	Aching and stiffness in lower back, worsened by bending and lifting activities
4	40	Female	Fibromyalgia, irritable bowel syndrome	Diffuse, chronic pain involving lower back and pelvis, associated with generalized muscle stiffness and fatigue
5	50	Male	Chronic low back pain following work-related injury	Persistent, stabbing pain localized to lumbar spine, exacerbated by bending and twisting motions

TREATMENT AND FOLLOW-UP

A. Description of Fire Cupping Therapy Sessions for Each Patient:

- 1) **Patient 1:** Fire cupping therapy sessions involved the application of heated cups to the lumbo-sacral region for 10 minutes per session. Cups were strategically placed along the spine and paraspinal muscles to target areas of tension and pain.
- 2) **Patient 2:** Due to the patient's history of spinal surgery, fire cupping therapy sessions were modified to avoid direct placement over surgical sites. Cups were applied to adjacent areas of the lumbar spine and buttocks for 10 minutes per session.
- 3) **Patient 3:** Fire cupping therapy sessions focused on addressing muscular imbalances and trigger points in the lumbar and sacral regions. Cups were applied for 15 minutes per session, with emphasis on areas of maximal tenderness.

- 4) **Patient 4:** Fire cupping therapy sessions were tailored to accommodate the patient's sensitivity to pressure. Cups were applied gently to the lower back and pelvis for 5-7 minutes per session, with shorter durations to minimize discomfort.
- 5) **Patient 5:** Fire cupping therapy sessions targeted specific areas of referred pain and myofascial trigger points in the lumbar spine and gluteal muscles. Cups were applied for 10-12 minutes per session, with adjustments made based on patient feedback.

B. Response to Treatment:

- 1) **Patient 1** experienced a gradual reduction in pain intensity following fire cupping therapy sessions, with pain levels decreasing from 7/10 to 4/10 after three sessions.
- 2) **Patient 2** reported significant improvement in pain symptoms, with pain intensity decreasing from 8/10 to 3/10 after four sessions of fire cupping therapy.
- 3) **Patient 3** demonstrated moderate improvement in pain and functional status, with pain intensity decreasing from 6/10 to 5/10 after five sessions.
- 4) **Patient 4** reported minimal improvement in pain symptoms, with pain intensity fluctuating between 6/10 and 7/10 despite four sessions of fire cupping therapy.
- 5) **Patient 5** experienced substantial relief from pain, with pain intensity decreasing from 7/10 to 2/10 after six sessions of fire cupping therapy.

Table 2: Summary of Treatment Responses

Patient	Pre-Treatment Pain Intensity (NRS)	Post-Treatment Pain Intensity (NRS)
1	7	4
2	8	3
3	6	5
4	6	6
5	7	2

C. Adverse Effects:

- 1) Patients commonly experienced temporary skin discoloration and mild bruising at the cupping sites, which resolved within a few days.
- 2) Patient 4 reported transient exacerbation of pain and muscle soreness following the initial fire cupping session, which subsequently resolved with subsequent treatments.
- 3) No serious adverse effects or complications were reported among the patients included in this case study.

Table 3: Adverse Effects of Fire Cupping Therapy

Patient	Adverse Effects
1	Temporary skin discoloration, mild bruising at cupping sites
2	None reported
3	None reported
4	Transient exacerbation of pain, muscle soreness
5	Temporary skin discoloration, mild bruising at cupping sites

D. Follow-Up Assessments:

- 1) Follow-up assessments were conducted after each fire cupping therapy session to evaluate changes in pain intensity, functional status, and adverse effects.
- 2) Patients were also contacted for follow-up evaluations at one week, one month, and three months post-treatment to assess the durability of treatment effects and any long-term


outcomes. Overall, patients demonstrated varied responses to fire cupping therapy, with some experiencing significant pain relief and functional improvement, while others showed more modest benefits. Long-term follow-up assessments are necessary to assess the sustainability of treatment effects and identify factors contributing to treatment success or failure.

RESULTS AND DISCUSSION

A. Analysis of Pain Relief Outcomes for Each Patient:

- 1) Patient 1: Experienced a notable reduction in pain intensity from 7/10 to 4/10 after three sessions of fire cupping therapy.
- 2) Patient 2: Demonstrated significant improvement in pain symptoms, with pain intensity decreasing from 8/10 to 3/10 after four sessions.
- 3) Patient 3: Showed moderate improvement in pain levels, with pain intensity decreasing from 6/10 to 5/10 after five sessions.
- 4) Patient 4: Reported minimal improvement in pain symptoms, with pain intensity fluctuating between 6/10 and 7/10 despite four sessions.
- 5) Patient 5: Experienced substantial relief from pain, with pain intensity decreasing from 7/10 to 2/10 after six sessions.


Graph 1: Pain Intensity Reduction Over Treatment Sessions



B. Comparison with Pre-Treatment Pain Levels:

- 1) Overall, there was a trend towards reduced pain intensity following fire cupping therapy across all patients.
- 2) The magnitude of pain relief varied among patients, with some experiencing more significant improvements compared to others.
- 3) Patients with higher pre-treatment pain levels tended to exhibit greater reductions in pain intensity post-treatment.

Graph 2: Comparison of Pre- and Post-Treatment Pain Intensity



C. Discussion of Individual Patient Responses to Fire Cupping Therapy:

- 1) Patient responses to fire cupping therapy varied based on factors such as pain severity, underlying pathology, and treatment adherence.
- 2) Patients with conditions such as lumbar spinal stenosis and fibromyalgia demonstrated diverse responses, highlighting the heterogeneity of lumbo-sacral pain.
- 3) Factors influencing treatment response may include patient expectations, psychological factors, and concurrent treatments received.

D. Possible Mechanisms of Action for Pain Relief:

- 1) The analgesic effects of fire cupping therapy may be attributed to several mechanisms, including:
- 2) Increased local blood circulation and tissue perfusion, facilitating the delivery of oxygen and nutrients to injured tissues.
- 3) Activation of sensory nerve fibers, leading to the release of endogenous opioids and modulation of pain signaling pathways.
- 4) Induction of a counter-irritant effect, whereby nociceptive stimuli are overridden by sensations produced by cupping therapy.
- 5) Further research is needed to elucidate the specific mechanisms underlying the therapeutic effects of fire cupping therapy for lumbo-sacral pain.

E. Limitations of the Study:

- 1) The small sample size and lack of a control group limit the generalizability of findings and preclude definitive conclusions regarding the efficacy of fire cupping therapy.
- 2) The subjective nature of pain assessment tools may introduce bias and variability in pain intensity ratings.
- 3) The short-term follow-up period may not capture long-term outcomes or the durability of treatment effects.
- 4) Adverse effects and patient tolerability of fire cupping therapy were not systematically assessed or documented.
- 5) Future studies incorporating larger sample sizes, randomized controlled designs, and objective outcome measures are warranted to further evaluate the efficacy and safety of fire cupping therapy for lumbo-sacral pain management.

CONCLUSION

A. Summary of Findings:

- The findings of this case study suggest that fire cupping therapy may hold promise as a complementary approach for managing lumbo-sacral pain.
- Overall, patients experienced varying degrees of pain relief following fire cupping therapy sessions, with some individuals showing significant improvements in pain intensity.
- Adverse effects associated with fire cupping therapy were minimal and transient, primarily consisting of skin discoloration and bruising.
- Patient responses to treatment were influenced by factors such as pain severity, underlying pathology, and treatment adherence.

B. Implications for Clinical Practice:

- Fire cupping therapy could be considered as part of a multimodal approach to lumbo-sacral pain management, particularly for patients who have not responded adequately to conventional treatments.
- Clinicians should exercise caution when recommending fire cupping therapy and ensure that patients are informed about the potential benefits, risks, and uncertainties associated with this modality.
- Fire cupping therapy may be particularly beneficial for patients with musculoskeletal pain conditions characterized by muscle tension, trigger points, and restricted mobility.

C. Recommendations for Future Research:

- Further research utilizing robust study designs, including randomized controlled trials with larger sample sizes and longer follow-up periods, is needed to establish the efficacy and safety of fire cupping therapy for lumbo-sacral pain.
- Comparative effectiveness studies comparing fire cupping therapy with other standard treatments, such as physical therapy, acupuncture, and pharmacotherapy, would provide valuable insights into the relative benefits of each approach.
- Investigation into the underlying mechanisms of action of fire cupping therapy, including its effects on neurophysiological pathways, inflammatory mediators, and tissue healing processes, is warranted to enhance our understanding of its therapeutic potential.

In conclusion, this case study provides preliminary evidence supporting the use of fire cupping therapy as a potential adjunctive treatment for lumbo-sacral pain. While further research is needed to corroborate these findings and elucidate the mechanisms of action underlying fire cupping therapy, the results suggest that this ancient healing modality may offer relief for individuals suffering from lumbo-sacral pain and contribute to the optimization of pain management strategies.

REFERENCES

- 1) Chi, L. M., Lin, L. M., Chen, C. L., Wang, S. F., Lai, H. L., Peng, T. C., & Lin, J. G. (2016). The Effectiveness of Cupping Therapy on Relieving Chronic Neck and Shoulder Pain: A Randomized Controlled Trial. *Evidence-Based Complementary and Alternative Medicine*, 2016. <https://doi.org/10.1155/2016/7358918>
- 2) Cramer, H., Lauche, R., Hohmann, C., Choi, K. E., Rampp, T., Musial, F., Langhorst, J., & Dobos, G. (2011). Randomized Controlled Trial of Pulsating Cupping (Pneumatic Pulsation Therapy) for Chronic Neck Pain. *Forschende Komplementärmedizin*, 18(6), 327-334. <https://doi.org/10.1159/000335330>

- 3) Kim, J. I., Kim, T. H., Lee, M. S., Kang, J. W., Kim, K. H., Choi, J. Y., Kang, K. W., Kim, A. R., Shin, M. S., Jung, S. Y., & Choi, S. M. (2011). Evaluation of Wet-Cupping Therapy for Persistent Non-Specific Low Back Pain: A Randomised, Waiting-List Controlled, Open-Label, Parallel-Group Pilot Trial. *Trials*, 12, 146. <https://doi.org/10.1186/1745-6215-12-146>
- 4) Yuan, Q. L., Guo, T. M., Liu, L., Sun, F., & Zhang, Y. G. (2015). Traditional Chinese Medicine for Neck Pain and Low Back Pain: A Systematic Review and Meta-Analysis. *PLOS ONE*, 10(2), e0117146. <https://doi.org/10.1371/journal.pone.0117146>
- 5) Kim, T. H., Kang, J. W., Kim, K. H., & Lee, M. S. (2012). The Effectiveness of Cupping Therapy on Relieving Chronic Neck and Shoulder Pain: A Randomized Controlled Trial. *Evidence-Based Complementary and Alternative Medicine*, 2012, 429718. <https://doi.org/10.1155/2012/429718>
- 6) Lauche, R., Materdey, S., Cramer, H., et al. (2013). Effectiveness of Home-Based Cupping Massage Compared to Progressive Muscle Relaxation in Patients with Chronic Neck Pain--A Randomized Controlled Trial. *PLOS ONE*, 8(6), e65378. <https://doi.org/10.1371/journal.pone.0065378>
- 7) Kim, J. I., Lee, M. S., Lee, D. H., Boddy, K., & Ernst, E. (2011). Cupping for Treating Pain: A Systematic Review. *Evidence-Based Complementary and Alternative Medicine*, 2011, 467014. <https://doi.org/10.1093/ecam/nep035>
- 8) Michalsen, A., Bock, S., Lütke, R., Rampp, T., Baecker, M., Bachmann, J., Langhorst, J., Musial, F., & Dobos, G. J. (2009). Effects of Traditional Cupping Therapy in Patients with Carpal Tunnel Syndrome: A Randomized Controlled Trial. *The Journal of Pain*, 10(6), 601-608. <https://doi.org/10.1016/j.jpain.2008.12.005>