
A Comparative Study of Serum Calcium Levels in Pre-Eclampsia Versus Normotensive Pregnant Women

Mandara TP, Ashok Kumar K**, Ranjitha A***, Priyanka Dapper*****

Junior Resident, Professor*** Junior Resident**** Junior Resident*
Department of OBG, ESIC & PGIMSR, Rajajinagar, Bangalore, Karnataka

***Corresponding author**

Email: manaadarahello@gmail.com

ABSTRACT

Introduction: Preeclampsia is described as a pregnancy specific syndrome that can affect virtually every organ system. Proteinuria is an objective marker and reflects the system wide endothelial leak that characterizes the preeclampsia syndrome. Preeclampsia is a leading cause of maternal, fetal, neonatal and perinatal mortality and morbidity worldwide. In our study, the aim was to compare Serum calcium levels in Preeclamptic patients and normal pregnancies and to determine the role of serum calcium level in Preeclamptic patients and normal pregnant women.

Material and Methods: We conducted a prospective comparative study. The study was undertaken in 90 pregnant women. Data for the study was collected from 45 normotensive pregnant women (control group) and 45 Preeclamptic patients (study group) with more than 24 weeks of gestational age at ESIC-MC & PGIMSR Hospital, Bangalore. Between March 2021 to August 2022.

Results: The mean BMI of cases is 25.03 (S.D=2.65), the mean BMI of controls is 22.89 (S.D=2.18). Student t-test shows that the BMI differs significantly between the two groups ($p < 0.005$). The mean systolic BP of cases is 155.47 mm/Hg (S.D=8.26 mm/Hg) and diastolic BP is 96.67 mm/Hg (S.D=5.54 mm/Hg). The mean systolic BP of controls is 119.80 mm/hg (S.D=8.12 mm/Hg) and diastolic BP is 77.20 mm/Hg (S.D=4.37 mm/Hg). The difference was statistically significant ($p < 0.001$). The mean serum calcium level of Preeclamptic women (8.23 mg/dl S.D=0.76)is significantly lower($p < 0.001$)than in normotensive patients (9.22 mg/dl S.D=0.59 mg/dl).

Conclusion: This study concludes that serum calcium levels were significantly low in preeclamptic patients. Thus, monitoring serum calcium levels can be effective in preventing maternal and fetal morbidity, mortality and help to reach a favorable outcome in pregnancy.

Key words: Preeclampsia, Hypertension, BMI, Albuminuria, Serum Calcium.

INTRODUCTION

Preeclampsia is best described as a pregnancy specific syndrome that can affect virtually every organ system. Although preeclampsia is much more than simply gestational hypertension with proteinuria, appearance of proteinuria is an important diagnostic criterion. Thus, proteinuria is an objective marker and reflects the system wide endothelial leak that characterizes the preeclampsia syndrome [1]. Preeclampsia usually begins after 20 weeks of pregnancy in a woman whose blood pressure had been normal previously. It can lead to serious, even fatal, complications to both mother and baby. Preeclampsia is associated with an increased risk of placental abruption, preterm birth, fetal intrauterine growth restriction (IUGR), acute renal failure, cerebrovascular and cardiovascular complications, disseminated intravascular coagulation, and maternal death. Therefore, the need to provide an early diagnosis of Preeclampsia is vital [1].

The requirement of calcium in pregnancy is very high and if unmet leads to demineralization of maternal skeleton, stunted fetal growth, reduced mineralization of the fetal bones and hampers the secretion of calcium in breast-milk [2]. By the end of pregnancy, around 25-30 g of calcium is transferred to the fetal skeleton from the mother. The accumulation begins at the rate of 2-3 mg/day of calcium in the first trimester increasing to 250 mg/day by the end of third trimester³. Loss of calcium in breast milk during lactation is around 200-240 mg/day [4]. Approximately, mother loses 3-5% of maternal skeletal calcium content [5]. Initial studies on calcium in pregnancy led to the hypothesis that high calcium intake reduces the incidence of Pregnancy induced hypertension. However, it could not be established the actual reason behind this [6]. There are not many studies that determine the role of serum calcium level in Pregnancy from the Indian population. Therefore, this study aims to assess the levels of calcium in normal pregnant women and in those with preeclampsia. The period of pregnancy and lactation is a demanding phase for a woman when the calcium demand increases enormously. The fetus and the neonate tend to rely on the mother for calcium. The excessive demand may at times exceed the normal daily intake of the woman. There are specific changes in the body during pregnancy to meet these changes in requirements. The major adaptation is doubling the calcium absorption during pregnancy. This adaptation in mothers also influences the way other disorders of calcium and bone are presented.

MATERIAL AND METHODS

Study type: Cross sectional study

Study place: ESIC-PGIMSR, Rajajinagar, Bangalore

Study period: March 2021 to August 2022

Sample size Calculation

45 normal pregnant women, 45 women with preeclampsia, a combined total of size was 90. Sample size was calculated based on a previous study conducted by Rajendra Kumar *et al.*,⁴⁸ in which it was observed that mean serum calcium level in the pre-eclampsia group was 8.69 ± 0.66 . In the present study considering the mean difference of 0.8, power of 90% and alpha error of 5%.sample size was estimated to be 45 in each group. (Total 90).

$$\text{Formula: } N = \frac{(Z_{\alpha/2} + Z_{\beta})^2 \sigma^2}{\delta^2}$$

Inclusion Criteria

Patients who are willing to give consent. 2. Patients diagnosed with preeclampsia based on criteria - blood pressure $\geq 140/90$ mmHg on two separate occasions 6 hours apart, 1+ dipstick in two midstream urine samples collected 4 hours apart, urine protein creatinine ratio >0.3 mg/dl in more than 20 weeks period of gestation. 3. Asymptomatic pregnant women with gestational age more than 24 weeks of pregnancy with blood pressure less than 130/80 mm Hg. 4. Singleton pregnancy 5. Age 18-40 years.

Exclusion Criteria

Patients not willing to give informed consent 2. Chronic hypertension 3. Type 1 and 2 diabetes mellitus 4. Diagnosed case of renal disease, cardiovascular disease, liver disorder. 5. K/C/O APLA Syndrome 6. Multiple pregnancy

PROCEDURE

After obtaining approval and clearance from the institutional ethics committee, considering the inclusion and exclusion criteria, all pregnant women were recruited from Department of Obstetrics and Gynecology, ESICMC & PGIMSR from MARCH 2021 to AUGUST 2022.


STATISTICAL ANALYSIS

- 1) Data recorded was statistically analysed. Data were entered in a predesigned proforma. (Annexure III) Quantitative variables like age, BMI, Blood pressure levels, calcium levels, urine protein creatinine ratio were analysed using descriptive statistics (mean + SD) or median (IQR). Categorical variables were presented using frequency and percentage. Continuous data were represented as mean and standard deviation and the p-value was calculated using an independent t-test.
- 2) The mean serum calcium levels (continuous variable) were compared between the 2 groups using student t-test.
- 3) Chi-square test was used as a test of significance for the association hypothesis for categorical/ qualitative data.
- 4) Pearson's correlation was used to assess the correlation between continuous variables and spearman correlation was used for categorical correlations.
- 5) Graphical representation of data: MS Excel and MS word was used to obtain the various bar diagrams. SPSS was used to generate the scatterplots for correlation.
- 6) p value (Probability that the result is not by chance) of <0.05 was considered as statistically significant after assuming all the rules of statistical tests.

RESULTS


In our study, the mean age of cases was 27.4 years (S.D=4.67 years) and mean age of controls was 25.53 years (S.D=4.64 years). Student t-test showed that the age does not differ significantly between the two groups ($p > 0.05$). Among the cases, 20 (44.4%) were primigravida, 25 (55.6%) was multigravida, while 25 (55.6%) of the controls were primigravida and 20 (44.44%) were multigravida. Chi-square analysis shows that the difference was not statistically significant ($\chi^2 = 7.510$; $p = 0.057$). The mean gestational age of cases was 36.69 ± 1.59 weeks and controls were 38.60 ± 1.25 weeks. Comparison of gestational age between cases and controls showed that there was statistically significant difference ($t = -6.33$; $p < 0.0001$).

Graph 1: Comparison of gestational age between cases and controls




The mean BMI of cases was 25.03 ± 2.65 Kg/m² and that of the controls was 22.89 ± 2.18 Kg/m². Student t-test showed that the mean difference of BMI between cases and controls was 2.14 Kg/m² and that this difference was statistically significant ($t=4.18$; $p < 0.05$) while only 7 controls were preobese. Overall, only 12 cases had normal BMI whereas 28 controls had normal prepregnancy BMIs. The remaining 33 cases were overweight (12), pre-obese (20) or obese (1).

Graph 2: Comparison of BMI between cases and controls




Graph 3: Comparison of Pre-pregnancy BMI between cases and controls using Asia-Pacific population




The mean systolic BP of cases was 155.47 ± 8.26 mmHg. The range was between 141 mmHg and 177 mm/Hg. The mean systolic BP of controls was 119.80 ± 8.12 mmHg. The range was between 101 mm/Hg and 135 mm/Hg. The mean diastolic BP of cases was 96.67 ± 5.54 mmHg. The range was between 83 mm/Hg and 106 mm/Hg. The mean diastolic BP of controls was 77.20 ± 4.37 mmHg. The range was between 71 mm/Hg and 86 mm/Hg. As the cases were by definition having elevated blood pressure, both SBP and DBP among cases were significantly higher than those of the controls, statistically significant ($p < 0.001$).

Graph 4: Comparison of Systolic BP between cases and controls




Graph 5: Comparison of Diastolic BP between cases and controls




The mean serum calcium level of cases was 8.23 ± 0.76 mg/dL (range 6.3 - 9.6 mg/dL). The mean serum calcium level of controls was 9.22 ± 0.59 mg/dL (range 8.4-10.6 mg/dL). A comparison of mean serum calcium levels between cases and controls shows that the mean serum calcium is lower by 0.99 mg/dL in cases. The difference was statistically significant ($p < 0.001$).

Graph 6: Comparison of serum calcium levels between cases and controls




Among the cases and controls, Serum Calcium levels were negatively correlated with Systolic Blood Pressure (Pearson Rho= -0.75). This strong correlation was statistically significant (p value.<0.001) Among the cases the correlation coefficient for Pearson’s correlation was -0.663 among the controls, it was -0.624. (both p values <0.0001). Among the cases and controls, Serum Calcium levels were negatively correlated with Diastolic Blood Pressure (Pearson Correlation Coefficient= -0.6). This correlation was statistically significant (p value<0.001). Among the cases the correlation coefficient for Pearson’s correlation was -0.502 (p value <0.001) among the controls, it was -0.39 (p-values=0.008).

Graph 7: Correlation between Serum Calcium and Systolic Blood Pressure




Graph 8: Correlation between Serum Calcium and Diastolic Blood Pressure



Urine PCR was found to be negatively correlated with Serum Calcium, but this was only among the cases (Pearson Correlation Coefficient= -0.691, p-value<0.0001); whereas among the controls there was no significant correlation (Pearson Correlation Coefficient= 0.147, p-value=0.335). This is to be expected as we do not usually see proteinuria among normal pregnant women (without pre-eclampsia).

Graph 9: Comparing urine PCR (protein creatinine ratio) with Serum Calcium



DISCUSSION

Our study aimed to determine the role of serum calcium level in Preeclampsia and normal pregnant women and to compare Serum calcium levels in Preeclamptic patients and normal pregnant women.

In our study, the mean age of cases was 27.4 years (S.D=4.67 years). It ranges between 21 and 38 years. The mean age of controls was 25.53 years (S.D=4.64 years). the age does not differ significantly between the two groups (p>0.05). Similarly no statistical significance were

noted in age between two groups in study conducted by Soumi B et al⁸, Leila F et al [9] the mean gestational age of cases was 36.69 weeks (S.D=1.59 weeks). The mean gestational age of controls was 38.60 months (S.D=1.25 weeks). Comparison of gestational age between cases and controls shows that there was statistically significant difference ($p < 0.001$). There was increased chance of preterm deliveries in preeclamptic patients, due to early termination because of complications arising from preeclampsia.

Our study is in line with study conducted by Endalamaw et al [10]. In This study the mean of age of cases is 35.68 ± 4.57 were and that of controls is 38.7 ± 3.19 and the difference is statistically significant ($p < 0.001$). Similar findings were noted in study conducted by Rahel D et al [11].

The mean BMI of cases was 25.03 ± 2.65 Kg/m² and that of the controls was 22.89 ± 2.18 Kg/m². Student t-test shows that the mean difference of BMI between cases and controls was 2.14 Kg/m² and that this difference was statistically significant ($t=4.18$; $p < 0.0001$). Pregnant women with higher BMI tend to develop preeclampsia. Similar findings were reported by Reetu R et al¹², Brandie D et al [13], and Laaya H et al [14].

The mean systolic BP of cases was 155.47 mm/Hg (S.D=8.26 mm/Hg). The range was between 141 mm/Hg and 177 mm/Hg. The mean systolic BP of controls was 119.80 mm/Hg (S.D=8.12 mm/Hg). The range was between 101 mm/Hg and 135 mm/Hg. Comparison of systolic BP between cases and controls showed that the mean systolic BP was higher in cases. The difference is statistically significant ($p < 0.005$).

The mean diastolic BP of cases was 96.67 mm/Hg (S.D=5.54 mm/Hg). The range was between 83 mm/Hg and 106 mm/Hg. The mean diastolic BP of controls was 77.20 mm/hg (S.D=4.37 mm/Hg). The range was between 71 mm/Hg and 86 mm/Hg. The median diastolic BP of controls was 74.22 mm/Hg. Comparison of diastolic BP between cases and controls showed that the mean diastolic BP is higher in cases. The difference is statistically significant ($p < 0.005$).

Thus, the findings in this study is in line with the findings of the study by Mohammed G et al [15], Ebenezer O et al [16], Deepa V et al [17], and Oneyama A et al [18].

The mean serum calcium level of cases was 8.23 ± 0.76 mg/dl. It ranges between 6.3-9.6 mg/dl. The mean serum calcium level of controls was 9.22 ± 0.59 . It ranges between 8.4-10.6 mg/dl. Comparison of mean serum calcium levels between cases and controls shows that the mean serum calcium was lower in cases. The difference was statistically significant ($p < 0.005$).

Our study is in line with a study conducted by Salonee M et al [19], in this study the mean serum calcium level in the cases was 8.33 ± 0.58 mg/dl and 8.93 ± 0.45 mg/dl in the controls. Comparison of the serum calcium (mg/dL) between the two groups shows that serum calcium (mg/dL) is higher in control group with a t value of 4.491 and is statistically significant.

Another meta-analysis of randomized control trials to understand the relationship between calcium supplementation, blood pressure, preeclampsia, and outcomes of pregnancy was done by Heiner et al in 1996²⁰. Fourteen trials were chosen and 2459 women were selected. The findings were differential and concluded that; Low calcium levels are associated with higher incidence of hypertension in pregnancy, Calcium supplementation reduced the systolic and diastolic blood pressure and Women at risk of pre-eclampsia can be advised calcium

supplementation.

Among the cases and controls, Serum Calcium levels were negatively correlated with Systolic Blood Pressure (Pearson Rho= -0.75). This strong correlation was statistically significant (p value<0.0001). Among the cases the correlation coefficient for Pearson's correlation was -0.663 among the controls, it was -0.624. (Both p values <0.0001).

Among the cases and controls, Serum Calcium levels were negatively correlated with Diastolic Blood Pressure (Pearson Correlation Coefficient= -0.6). This correlation was statistically significant (p value<0.0001). Among the cases the correlation coefficient for Pearson's correlation was -0.502 (p-value<0.0001) among the controls, it was -0.39 (p-values=0.008).

Findings in our study was in line with study conducted by Sultana P et al [21], In this study the mean serum calcium was 8.2 ± 0.2 mg/dl in Group A, 7.6 ± 0.3 mg/dl in Group B and 9.0 ± 0.5 mg/dl in Group C. The difference was statistically significant (p<0.005) among three groups. The mean serum calcium was significantly lower in severe preeclampsia as compared to mild preeclampsia and normal pregnant women.

Pearson's correlation test revealed that both systolic blood pressure (r= -0.746; p=0.001) and diastolic blood pressure (r= -0.688; p=0.001) had negative correlation with the serum calcium concentration.

Thus, in our study Both the systolic and diastolic blood pressure had negative association with low serum calcium concentration.

In our study, the mean BMI, Serum Calcium, Systolic BP, and Diastolic BP differs significantly between the two groups. There is a reduction in the serum calcium levels (0.99mg/dL) of cases of Pre-eclampsia as compared to the normal pregnant women.

Therefore, there is a definite relationship between serum calcium levels and hypertensive disorders of pregnancy. This study shows a relationship between serum calcium levels and blood pressure of the pregnant women. The primary event leading to hypertension in preeclampsia is unknown. Studies show that before the clinical presentation of the disease, the pathophysiological changes are seen. This study concludes that regular monitoring of patients for these biomarkers will aid in early diagnosis and management. Thus, monitoring serum calcium levels and supplementing with calcium can be effective in preventing maternal morbidity and mortality and help to reach a favourable outcome in pregnancy.

CONCLUSION

The result of the present study suggests that hypocalcaemia was present in preeclamptic pregnant women and serum calcium was significantly lower than in normal pregnant women. Thus, use of serum calcium may help to identify pregnant women at risk of preeclampsia and prompt initiation of education and prophylactic interventions.

REFERENCES

- 1) Bloom SL, Dashe JS, Hoffman BL, Spong CY..Hypertensive disorders Williams obstetrics,25th ed. New York:McGraw-Hill Education Medical;2022.p.711-12
- 2) Prentice, A. (2000). Calcium in pregnancy and lactation. *Annual review of nutrition*,

- 20(1), 249-272.
- 3) Widdowson, E. M. (1981). Changes in body composition during growth. *Scientific foundations of paediatrics*, 330-342.
 - 4) Laskey, M. A., Prentice, A., Hanratty, L. A., Jarjou, L. M., Dibba, B., Beavan, S. R., & Cole, T. J. (1998). Bone changes after 3 mo of lactation: influence of calcium intake, breast-milk output, and vitamin D-receptor genotype. *The American journal of clinical nutrition*, 67(4), 685-692.
 - 5) Kalkwarf, H. J. (2006). Calcium in pregnancy and lactation. In *Calcium in Human Health* (pp. 297-309). Humana Press.
 - 6) Belizan, J. M., & Villar, J. (1980). The relationship between calcium intake and edema-, proteinuria-, and hypertension-gestosis: A hypothesis. *The American journal of clinical nutrition*, 33(10), 2202-2210.
 - 7) Chaudhari RK, Niraula A, Bataju M, Baranwa JK, Khan SA, Bhatt RD, Chettr S, Lamsa M. Serum Calcium and Magnesium levels in Pre-eclampsia. *National Journal of Laboratory Medicine*. 2018.
 - 8) Biswas S, Roy A, Biswas S. Comparative study of copper, zinc, iron, ferritin, calcium and magnesium levels in pregnancy induced hypertension and normotensive primigravida mothers. *Int. J. Res. Med. Sci.* 2016 Jun;4:1879-83.
 - 9) Farzin L, Sajadi F. Comparison of serum trace element levels in patients with or without pre-eclampsia. *Journal of research in medical sciences: the official journal of Isfahan University of Medical Sciences*. 2012 Oct;17(10):938.
 - 10) Tesfa E, Munshea A, Nibret E, Mekonnen D, Sinishaw MA, Gizaw ST. Evaluation of maternal serum lipid profiles and clinical chemistry parameters in the prediction of pre-eclampsia in pregnant women attending ANC and delivery services. (2023)
 - 11) Gebreyohannes RD, Abdella A, Ayele W, Eke AC. Association of dietary calcium intake, total and ionized serum calcium levels with preeclampsia in Ethiopia. *BMC pregnancy and childbirth*. 2021 Dec;21(1):1-7.
 - 12) Rani R, Kumar S, Singh SK. Study of Serum Calcium and Magnesium levels in Pre-eclampsia at DMC, Laheriasarai, Bihar. (2022)
 - 13) Taylor BD, Ness RB, Olsen J, Hougaard DM, Skogstrand K, Roberts JM, Haggerty CL. Serum leptin measured in early pregnancy is higher in women with preeclampsia compared with normotensive pregnant women. *Hypertension*. 2015 Mar;65(3):594-9.
 - 14) Hamedanian L, Badehnoosh B, Razavi-Khorasani N, Mohammadpour Z, Mozaffari-Khosravi H. Evaluation of vitamin D status, parathyroid hormone, and calcium among Iranian pregnant women with preeclampsia: A case-control study. *International Journal of Reproductive BioMedicine*. 2019 Nov;17(11):831.
 - 15) Gamal Mohammed El-Maghraby M, Mohammed Zakarya AE, Abd EL-Latif Hashish M, Aly El-Boghdady A. Comparative Study Between Serum Calcium and Magnesium Levels in Pre-Eclampsia Versus Normal Pregnancy. *Al-Azhar Medical Journal*. 2022 Apr 1;51(2):999-1014.
 - 16) Darkwa EO, Antwi-Boasiako C, Djagbletey R, Owoo C, Obed S, Sottie D. Serum magnesium and calcium in preeclampsia: a comparative study at the Korle-Bu Teaching Hospital, Ghana. *Integrated blood pressure control*. 2017;10:9.
 - 17) Kanagal DV, Rajesh A, Rao K, Devi UH, Shetty H, Kumari S, Shetty PK. Levels of serum calcium and magnesium in pre-eclamptic and normal pregnancy: A study from Coastal India. *Journal of clinical and diagnostic research: JCDR*. 2014 Jul;8(7):OC01.
 - 18) Onyegbule OA, Meludu SC, Dioka CE, Udigwe GO, Udo JN, Ezidigboh AN, Osakue N. Comparison of serum levels of calcium and magnesium among preeclamptic and normotensive pregnant women at Nnamdi Azikiwe University Teaching Hospital, Nnewi, Nigeria. *Int J Res Med Sci*. 2014 Apr;2(2):404-8.

- 19) Naik SM, Sehra R, Bansal M. Association of serum calcium and serum magnesium in gestational hypertension and pre-eclampsia. *International Journal of Clinical Obstetrics and Gynaecology*. 2022;6(4):07-11.
- 20) Bucher HC, Guyatt GH, Cook RJ, Hatala R, Cook DJ, Lang JD, Hunt D. Effect of calcium supplementation on pregnancy-induced hypertension and preeclampsia: a meta-analysis of randomized controlled trials. *Jama*. 1996 Apr 10;275(14):1113-7.
- 21) Parvin S, Chowdhury SB, Nahar K, Hoque MM. Serum Calcium and Its Association with Preeclampsia. *Bangladesh J Med Sci [Internet]*. 2021 Feb. 1 [cited 2023 Jan. 15];20(2):379-83.